

COGNITIVE NEUROSCIENCE

A single dose of lorazepam reduces paired-pulse suppression of median nerve evoked somatosensory evoked potentials

Philipp Stude,¹ Melanie Lenz,¹ Oliver Höffken,¹ Martin Tegenthoff¹ and Hubert Dinse^{1,2}¹Department of Neurology, BG-Universitätsklinikum Bergmannsheil Bochum, Buerkle-de-la-Camp-Platz 1, 44789 Bochum, Germany²Neural Plasticity Laboratory, Institute for Neuroinformatics, Ruhr University, Bochum, Germany

Keywords: cortical excitability, GABA, intracortical inhibition, paired-pulse behaviour, plasticity, somatosensory evoked potentials

Edited by John Foxe

Received 28 April 2015, revised 14 February 2016, accepted 23 February 2016

Abstract

Paired-pulse behaviour in the somatosensory cortex is an approach to obtain insights into cortical processing modes and to obtain markers of changes of cortical excitability attributable to learning or pathological states. Numerous studies have demonstrated suppression of the response to the stimulus that follows a first one after a short interval, but the underlying mechanisms remain elusive, although there is agreement that GABAergic mechanisms seem to play a crucial role. We therefore aimed to explore the influence of the GABA_A agonist lorazepam on paired-pulse somatosensory evoked potentials (SEPs). We recorded and analysed SEPs after paired median nerve stimulation in healthy individuals before and after they had received a single dose of 2.5 mg of lorazepam as compared with a control group receiving placebo. Paired-pulse suppression was expressed as a ratio of the amplitudes of the second and the first peaks. We found that, after lorazepam application, paired-pulse suppression of the cortical N20 component remained unchanged, but suppression of the N20–P25 complex was significantly reduced, indicative of GABAergic involvement in intracortical processing. Our data suggest that lorazepam most likely enhances inhibition within the cortical network of interneurons responsible for creating paired-pulse suppression, leading to reduced inhibitory drive with a subsequently reduced amount of suppression. The results provide further evidence that GABA_A-mediated mechanisms are involved in the generation of median nerve evoked paired-pulse suppression.

Introduction

Stimulation with pairs of stimuli in close succession (paired-pulse stimulation) has become a common tool for investigating paired-pulse suppression (PPS). PPS describes the phenomenon whereby, at short interstimulus intervals (ISIs), cortical responses to the second stimulus are significantly reduced. PPS is quantified in terms of the amplitude of the second response divided by the amplitude of the first response. Accordingly, small amplitude ratios are associated with strong PPS, and large amplitude ratios are associated with reduced PPS. For the somatosensory system, PPS in combination with somatosensory evoked potential (SEP) recordings over the primary somatosensory cortex has been increasingly used to investigate paired-pulse behaviour, in order to obtain insights into cortical processing modes and to obtain markers of changes of cortical excitability attributable to learning or pathological states (Allison, 1962; Schwartz & Shagass, 1964; Shagass & Schwartz, 1964;

Ragert *et al.*, 2004; Höffken *et al.*, 2007, 2013a,b; Lenz *et al.*, 2011; Gatica Tossi *et al.*, 2013).

Despite substantial experimental and theoretical work, the mechanisms mediating paired-pulse behaviour are not fully understood. Because of differences in PPS between cortical and thalamic cells, it has been argued that inheritance of thalamic response properties is unlikely to account for long-lasting forward suppression (Wehr & Zador, 2005). For human subjects, on the basis of multichannel SEP recordings after paired median nerve stimulation, it has been shown that PPS is generated at least rostral to the brainstem nuclei (Höffken *et al.*, 2010). There is agreement that presynaptic mechanisms play a crucial role (Hashimoto & Kano, 1998). Wehr & Zador (2005) reported that, in the rat auditory cortex, GABA receptor-mediated inhibition does not play a major role in forward suppression for ISIs of < 100 ms. For longer ISIs, synaptic depression is assumed to be responsible for the observed PPS (Wehr & Zador, 2005). In the visual cortex, suppression is also more consistent with thalamocortical synaptic depression than with inhibition (Carandini *et al.*, 2002; Freeman *et al.*, 2002). There is also evidence for the

Correspondence: Dr P. Stude, as above.
E-mail: philipp.stude@bergmannsheil.de

involvement of glutamatergic transmission in the paired-pulse phenomenon (Takahashi *et al.*, 1996; von Gersdorff *et al.*, 1997).

In the motor domain, paired-pulse transcranial magnetic stimulation has been widely used to study intracortical inhibition of the human motor cortex. These studies provided several lines of evidence for a critical role of GABAergic, presumably GABA_A-mediated, inhibition (Kujirai *et al.*, 1993; Ziemann *et al.*, 1996, 2001; Werhahn *et al.*, 1999; Hanajima *et al.*, 2003; Florian *et al.*, 2008), although the involvement of GABA_B has also been advocated (Porter & Nieves, 2004), owing to pharmacological interventions (Florian *et al.*, 2008) or based on the timing of long-interval intracortical inhibition (Fitzgerald *et al.*, 2009).

Given the overwhelming evidence from motor cortex studies for a critical role of GABA_A, we sought to revisit the GABA_A influence on PPS in the primary somatosensory cortex of healthy adults. We found that, after lorazepam application, PPS of the cortical N20 component remained unchanged, but suppression of the N20–P25 complex was significantly reduced, indicative of GABAergic involvement in intracortical processing.

Materials and methods

Subjects

We investigated two groups of right-handed subjects: the target group (lorazepam) consisted of 13 subjects (six females and seven males; mean age, 24.6 years; standard deviation, ± 2.5 years); the control group (placebo) consisted of 10 subjects (five females and five males; mean age, 25.9 years; standard deviation, ± 3.6 years). All subjects underwent clinical neurological investigations to exclude somatic illness before their participation, and gave their written informed consent. The study was approved by the local Ethics Committee, and was performed in accordance with the Declaration of Helsinki.

Paired-pulse stimulation

To study changes in PPS, we applied a paired-pulse protocol described in a recent review and systematic analysis by Höffken *et al.* (2013a,b). It consists of paired electrical median nerve stimulation with an ISI of 30 ms. Nerve stimulation of the right side was performed with a block electrode on the wrist (pulse width of rectangular pulse of 0.2 ms; repetition rate of paired stimuli of 2 Hz). To verify the correct positioning of the stimulation electrode, subjects had to report a prickling sensation in the thumb, index, and middle finger. The stimulation intensity was chosen at 2.5-fold of the somatosensory threshold, and was kept constant for each subject before and after administration of drug. In all subjects, the stimulation intensity evoked a small twitch of the thenar muscles. During stimulation, subjects were seated in a comfortable chair and were instructed to relax but to stay awake with eyes closed.

Electroencephalography signals were recorded continuously with Ag–AgCl electrodes (resistance, < 5 k Ω) between C3' and Fz as the reference. C3' is located over the left primary somatosensory cortex (SI), 2 cm posterior to C3, according to the 10/20 system. The electroencephalography signals were amplified with a bandwidth of 0.1–1000 Hz and digitized at 2.5 kHz with the BrainAmp Amplifier (Brainproducts, Munich, Germany). The electrical potentials were segmented in epochs from -50 ms to 200 ms, baseline-corrected, and averaged. Latencies and peak-to-peak amplitudes of the cortical N20 and the N20–P25 response components were compared before and after drug administration. PPS was calculated as a ratio of the

amplitude of the second response peak (A2) and the amplitude of the first response peak (A1), i.e. A2/A1 (Fig. 1). Analysis was performed by a blinded coworker who was not involved in data recording, using BRAINVISION ANALYZER (Brainproducts). A repeated-measures ANOVA with the within-subjects factor 'course: before/after' and the between-subjects factor 'group' (lorazepam group vs. control group) was performed.

When repeated-measures ANOVAs were used, all *F*-ratios associated with the repeated-measures factors were assessed by the use of degrees of freedom corrected with the Wilks' lambda procedure for controlling type I error. All statistical analyses were performed with SPSS 17 (SPSS, Chicago, IL, USA).

Electroencephalography recordings were performed in two sessions: the first session was used to obtain baseline data (before), and the second session started 75 min after intake of lorazepam (after). Each session contained a total number of 1000 paired-pulse stimuli. Medication consisted of a waver either of placebo or of 2.5 mg of lorazepam, and was administered in a pseudorandomized and double-blinded manner. Immediately after the end of the recording, blood samples were taken to quantify the plasma level of lorazepam.

Results

In repeated-measures ANOVA with the within-groups factor before–after and the between-subjects factor group (lorazepam group vs. control group), analysis of latencies and peak-to-peak amplitudes of both the N20 and the N20–P25 components revealed no significant change between the groups before and after lorazepam intake ($F = 1.757$, $P = 0.179$; Table 1). In the placebo group, the




FIG. 1. Single-subject SEPs following paired stimulation before (top) and after (bottom) intake of 2.5 mg of lorazepam. The amplitudes of the N20–P25 complex (A1 and A2) are marked.

TABLE 1. Amplitude and excitability parameters (means and standard errors of the mean) and *P*-values of *t*-tests of differences after administration of placebo or lorazepam

Parameter	Group	Mean	SD	Unit	<i>t</i> -value	<i>P</i> -value
A2/A1	Placebo	-0.057	0.079		-2.879	0.009
	Lorazepam	0.220	0.056			
A1	Placebo	-0.146	0.415	μV	0.911	0.373
	Lorazepam	-0.595	0.277	μV		
A2	Placebo	-0.130	0.147	μV	-1.258	0.223
	Lorazepam	0.194	0.193	μV		
N20*	Placebo	-0.013	0.238	μV	0.651	0.522
	Lorazepam	-0.232	0.226	μV		
N20†	Placebo	-0.121	0.315	μV	0.584	0.566
	Lorazepam	-0.324	0.173	μV		
P25*	Placebo	0.192	0.350	μV	0.832	0.415
	Lorazepam	-0.179	0.274	μV		
P25†	Placebo	-0.235	0.592	μV	0.075	0.941
	Lorazepam	-0.286	0.360	μV		

A1, amplitude of the first response peak; A2, amplitude of the second response peak; SD, standard deviation.

*Components of the SEP after the first of the paired stimuli.

†Components of the SEP after the second of the paired stimuli.

amplitudes of the first response of the N20 component were 1.76 ± 0.79 mV before intake and 1.75 ± 0.76 mV after intake, and the amplitudes of the second response were 1.53 ± 0.89 mV before intake and 1.41 ± 0.66 mV after intake (*t*-test: $P = 0.95$ and $P = 0.71$, respectively). Similarly, the amplitudes of the first and second N20–P25 components did not differ significantly before intake (2.86 ± 1.06 mV and 1.64 ± 0.81 mV, respectively) and after intake (2.72 ± 0.78 mV and 1.51 ± 0.71 mV, respectively) (*t*-test: $P = 0.73$ and $P = 0.4$, respectively).

A similar observation was made for the lorazepam group. For the N20 component, there was no significant difference between the before and after sessions for the first amplitude (1.51 ± 0.97 mV and 1.27 ± 0.69 mV, respectively; *t*-test, $P = 0.34$) or the second amplitude (1.29 ± 0.58 mV and 0.87 ± 0.51 mV, respectively; *t*-test, $P = 0.11$). For the N20–P25 component, neither the first response amplitudes (3.14 ± 1.36 mV before intake, and 2.54 ± 1.07 mV after intake; *t*-test, $P = 0.09$) nor the second response amplitudes (1.84 ± 0.64 mV before intake, and 2.04 ± 0.91 mV after intake) differed between the before and after sessions (*t*-test, $P = 0.35$). In contrast, ANOVA showed a significant effect for the within-subjects factor before–after \times group ($F = 3.097$, $P = 0.038$). Although no significant change in the A2/A1 ratio was found in the placebo group (0.61 ± 0.28 and 0.55 ± 0.28 , $P = 0.49$), we found a significant increase in the lorazepam group (0.65 ± 0.24 and 0.87 ± 0.33 ; *t*-test, $P = 0.003$), indicating that PPS was decreased after intake of lorazepam. This increase in the A2/A1 ratio after lorazepam intake was also significantly different from what was seen in the placebo group (0.55 ± 0.28 vs. 0.87 ± 0.33 ; *t*-test, $P = 0.026$; Fig. 2; Table 1).

The plasma level of lorazepam ranged between 16 ng/mL and 127 ng/mL. There was no correlation between concentration and electrophysiological parameters.

Discussion

Our study addressed the putative role of GABAergic transmission in the generation of paired-pulse behaviour in the somatosensory cortex. Using paired median nerve stimulation to record SEPs in the human somatosensory cortex, we demonstrated that a single dose of




FIG. 2. Mean paired-pulse ratios before (white) and after (light grey) lorazepam administration for the placebo group (left) and the lorazepam group (right). Error bars show the standard errors of the mean. Note the significant increase in the A2/A1 ratio in the lorazepam group ($***P = 0.003$), which was also significantly different from placebo ($*P = 0.026$).

the GABA_A agonist lorazepam modulates PPS of the cortical N20–P25 component, but not of the N20 component. Although numerous studies have demonstrated suppression of a response to a stimulus that follows a first one after a short interval, the underlying mechanisms remain elusive. There is agreement that PPS is most likely an intracortical phenomenon that is not present in a comparable form at a downstream level. Also, because of the suppressive nature of the paired-pulse behaviour, GABAergic contributions were advocated early on. However, data from animal research that have allowed detailed analysis of cellular and synaptic processes in mediating paired-pulse behaviour have created a complex picture (Wehr & Zador, 2005).

In contrast, studies in human individuals have to rely on more indirect approaches. For many years, paired-pulse transcranial magnetic stimulation has been widely used to study intracortical inhibition of the human motor cortex. The combination of paired-pulse transcranial magnetic stimulation with the application of agonists and/or antagonists of well-known transmitter systems has generated a considerable amount of information that, indeed, supports a role of GABAergic mechanisms in the generation of PPS (Kujirai *et al.*, 1993; Ziemann *et al.*, 1996, 2001; Werhahn *et al.*, 1999; Hanajima *et al.*, 2003; Di Lazzaro *et al.*, 2006; Florian *et al.*, 2008). Most studies have supported the involvement of GABA_A, but have also demonstrated that different forms of intracortical inhibition, such as short-interval intracortical inhibition, long-interval intracortical inhibition, and short-interval interhemispheric inhibition, might be mediated by different mechanisms (Florian *et al.*, 2008). PPS has been documented for the motor cortex (Ziemann *et al.*, 1996, 2001; Werhahn *et al.*, 1999; Hanajima *et al.*, 2003; Di Lazzaro *et al.*, 2006; Florian *et al.*, 2008), the auditory cortex (Percaccio *et al.*, 2005; Wehr & Zador, 2005), the somatosensory cortex (Allison, 1962; Schwartz & Shagass, 1964; Shagass & Schwartz, 1964; Ragert *et al.*, 2004; Höffken *et al.*, 2007; Lenz *et al.*, 2011), the visual cortex (Musselwhite & Jeffreys, 1983; Höffken *et al.*, 2008, 2009, 2013a,b), and higher cortical areas such as the dorsolateral prefrontal cortex (Fitzgerald *et al.*, 2009), implying that PPS is a ubiquitous cortical phenomenon that is not limited to a particular area. On the other hand, there are also significant differences in the properties of PPS across areas and modalities. For the somatosensory cortex, paired median nerve stimulation creates significant suppression up to

100 ms, whereas, for the visual cortex, suppression has been demonstrated for up to 200 ms or more (Höffken *et al.*, 2008). In contrast, for the motor cortex, transcranial magnetic stimulation-induced suppression evokes many forms of inhibition, such as short-interval intracortical inhibition and long-interval intracortical inhibition, which do not have an obvious equivalent in sensory cortices. In an early study of the effects of lorazepam on motor cortex excitability, corticocortical inhibition showed a tendency towards more inhibition, whereas corticocortical facilitation was almost completely suppressed (Ziemann *et al.*, 1996). In another study, in which GABAergic mechanisms were explored after application of the GABA uptake blocker tiagabine, PPS of the motor evoked potential at an ISI of 160 ms was more pronounced, whereas paired-pulse facilitation at an ISI of 10 ms was increased (Werhahn *et al.*, 1999).

In this and in our previous studies (Höffken *et al.*, 2007; Lenz *et al.*, 2012), we did not use a subtraction approach, but used raw amplitudes. In our view, the use of subtraction implicitly assumes that the response behaviour for single or paired stimulation is linear; that is, all of the later components present after single stimulation will show up identically under a paired-stimulation condition. In contrast, we took a more non-linear view, whereby new inputs, here the second stimulus, can override, reset or modulate components that are present under single stimulation. Therefore, in our view, the use of raw amplitudes involves fewer assumptions than does the subtraction approach. Moreover, in some of our previous studies, we had correlated paired-pulse ratios with perceptual measures, which revealed significant correlations, whereby the amount of PPS was linked to individual tactile performance behaviour (Höffken *et al.*, 2007; Lenz *et al.*, 2012). Such behaviour strongly supports the assumption that raw amplitude analysis provides a meaningful measure of intracortical excitability.

In our data, we found that, with an ISI of 30 ms, the response amplitudes of the first and second responses of the N20 and the N20–P25 components were not significantly altered. In contrast, the amplitude ratios of the N20–P25 component were significantly increased, indicative of reduced suppression. To explain these, on first glance, counterintuitive observations, we assume that the GABA_A agonist lorazepam most likely enhances inhibition within the network of interneurons responsible for creating PPS. As a result, the inhibitory drive is reduced, thereby reducing the amount of suppression. Interestingly, no such effects were observed for the N20 component of the SEPs. Further studies are needed to explore whether PPS evoked at ISIs shorter or longer than 30 ms shows a similar pharmacological dependency as demonstrated for 30 ms (Werhahn *et al.*, 1999; Wehr & Zador, 2005).

There is general agreement that the N20 component originates mainly in the granular layer (layer IV) of Brodmann's area 3b, which occupies the posterior bank of the Rolandic fissure (Allison *et al.*, 1989, 1991; McLaughlin & Kelly, 1993; Urbano *et al.*, 1997; Balzamo *et al.*, 2004). The origin of the P25 component is less clear. It has been proposed that the P25 component reflects the depolarization of the superficial portion of apical dendrites located in cortical layers 2 and 3 (Mitzdorf, 1985; Vaughan & Arezzo, 1988; Allison *et al.*, 1991; McLaughlin & Kelly, 1993; Nicholson Peterson *et al.*, 1995). Other studies have suggested a radially oriented source that is usually identified as Brodmann's area 1 at the apex of the postcentral gyrus (Arezzo *et al.*, 1979; Allison *et al.*, 1989, 1991; McCarthy *et al.*, 1991). Despite these discrepancies, there is agreement that the N20 component reflects thalamocortical input to SI, whereas the N20–P25 component represents intracortical processing (Wolters *et al.*, 2005). Such a dissociation is compatible

with our finding of a lack of GABAergic modulation of the N20 component. It should be noted that, owing to the cephalic channel recording used in this study, it is possible that the P25 potential is modulated by an N30 potential of frontal origin, further complicating the discussion of the origin of the P25 component.

By the use of magnetoencephalography to record somatosensory evoked magnetic fields, a GABAergic contribution has been studied following lorazepam administration (Huttunen *et al.*, 2008). These authors reported that, for ISIs of 20 ms, the drug had no effect on PPS or recovery for the N20 m deflection, but that the P35 m deflection was attenuated and did not recover at ISIs of 100 ms (Huttunen *et al.*, 2008). The lack of recovery at ISIs of 100 ms is difficult to reconcile with the early components seen in electrical SEPs, which all show recovery.

Although our data show clear involvement of GABA_A-mediated mechanisms in the generation of median nerve evoked PPS, research findings from the motor cortex and other areas make it highly likely that other mechanisms are also involved. For example, for the visual cortex, noradrenergic modulation of PPS has recently been demonstrated (Höffken *et al.*, 2012). Apparently, more studies are needed, including animal studies, to unravel the mechanisms of paired-pulse behaviour.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 874 A1 to M. Tegenthoff, M. Lenz and O. Höffken; SFB 874 A5 to H. Dinse; DFG Research Unit 1581 to M. Tegenthoff).

Abbreviations

A1, amplitude of the first response peak; A2, amplitude of the second response peak; ISI, interstimulus interval; PPS, paired-pulse suppression; SEP, somatosensory evoked potential.

References

- Allison, T. (1962) Recovery functions of somatosensory evoked responses in man. *Electroen. Clin. Neuro.*, **14**, 331–343.
- Allison, T., McCarthy, G., Wood, C.C., Darcey, T.M., Spencer, D.D. & Williamson, P.D. (1989) Human cortical potentials evoked by stimulation of the median nerve. I. Cytoarchitectonic areas generating short-latency activity. *J. Neurophysiol.*, **62**, 694–710.
- Allison, T., McCarthy, G., Wood, C.C. & Jones, S.J. (1991) Potentials evoked in human and monkey cerebral cortex by stimulation of the median nerve. A review of scalp and intracranial recordings. *Brain*, **114**, 2465–2503.
- Arezzo, J., Legatt, A.D. & Vaughan, H.G. Jr (1979) Topography and intracranial sources of somatosensory evoked potentials in the monkey. I. Early components. *Electroen. Clin. Neuro.*, **46**, 155–172.
- Balzamo, E., Marquis, P., Chauvel, P. & Régis, J. (2004) Short-latency components of evoked potentials to median nerve stimulation recorded by intracerebral electrodes in the human pre- and postcentral areas. *Clin. Neurophysiol.*, **115**, 1616–1623.
- Carandini, M., Heeger, D.J. & Senn, W. (2002) A synaptic explanation of suppression in visual cortex. *J. Neurosci.*, **22**, 10053–10065.
- Di Lazzaro, V., Pilato, F., Dileone, M., Ranieri, F., Ricci, V., Profice, P., Bria, P., Tonali, P.A. & Ziemann, U. (2006) GABAA receptor subtype specific enhancement of inhibition in human motor cortex. *J. Physiol.*, **575**, 721–726.
- Fitzgerald, P.B., Maller, J.J., Hoy, K., Farzan, F. & Daskalakis, Z.J. (2009) GABA and cortical inhibition in motor and non-motor regions using combined TMS-EEG: a time analysis. *Clin. Neurophysiol.*, **120**, 1706–1710.
- Florian, J., Müller-Dahlhaus, M., Liu, Y. & Ziemann, U. (2008) Inhibitory circuits and the nature of their interactions in the human motor cortex: a pharmacological TMS study. *J. Physiol.*, **586**, 495–514.
- Freeman, T.C., Durand, S., Kiper, D.C. & Carandini, M. (2002) Suppression without inhibition in visual cortex. *Neuron*, **35**, 759–771.

- Gatica Tossi, M.A., Lillemeier, A.S. & Dinse, H.R. (2013) Influence of stimulation intensity on paired-pulse suppression of human median nerve somatosensory evoked potentials. *NeuroReport*, **24**, 451–456.
- von Gersdorff, H., Schneggenburger, R., Weis, S. & Neher, E. (1997) Presynaptic depression at a calyx synapse: the small contribution of metabotropic glutamate receptors. *J. Neurosci.*, **17**, 8137–8146.
- Hanajima, R., Furubayashi, T., Iwata, N.K., Shiio, Y., Okabe, S., Kanazawa, I. & Ugawa, Y. (2003) Further evidence to support different mechanisms underlying intracortical inhibition of the motor cortex. *Exp. Brain Res.*, **151**, 427–434.
- Hashimoto, K. & Kano, M. (1998) Presynaptic origin of paired-pulse depression at climbing fibre–Purkinje cell synapses in the rat cerebellum. *J. Physiol.*, **506**, 391–405.
- Höfken, O., Veit, M., Knossalla, F., Lissek, S., Bliem, B., Ragert, P., Dinse, H.R. & Tegenthoff, M. (2007) Sustained increase of somatosensory cortex excitability by tactile coactivation studied by paired median nerve stimulation in humans correlates with perceptual gain. *J. Physiol.*, **584**, 463–471.
- Höfken, O., Grehl, T., Dinse, H.R., Tegenthoff, M. & Bach, M. (2008) Paired-pulse behavior of visually evoked potentials recorded in human visual cortex using patterned paired-pulse stimulation. *Exp. Brain Res.*, **188**, 427–435.
- Höfken, O., Stude, P., Lenz, M., Bach, M., Dinse, H.R. & Tegenthoff, M. (2009) Visual paired-pulse stimulation reveals enhanced visual cortex excitability in migraineurs. *Eur. J. Neurosci.*, **30**, 714–720.
- Höfken, O., Lenz, M., Tegenthoff, M. & Schwenkreis, P. (2010) Multichannel SEP-recording after paired median nerve stimulation suggests origin of paired-pulse inhibition rostral of the brainstem. *Neurosci. Lett.*, **468**, 308–311.
- Höfken, O., Lenz, M., Höckelmann, N., Dinse, H.R. & Tegenthoff, M. (2012) Noradrenergic modulation of human visual cortex excitability assessed by paired-pulse visual-evoked potentials. *NeuroReport*, **23**, 707–711.
- Höfken, O., Tannwitz, J., Lenz, M., Sczesny-Kaiser, M., Tegenthoff, M. & Schwenkreis, P. (2013a) Influence of parameter settings on paired-pulse-suppression in somatosensory evoked potentials: a systematic analysis. *Clin. Neurophysiol.*, **124**, 574–580.
- Höfken, O., Lenz, M., Sczesny-Kaiser, M., Dinse, H.R. & Tegenthoff, M. (2013b) Phosphene thresholds correlate with paired-pulse suppression of visually evoked potentials. *Brain Stimul.*, **6**, 118–121.
- Huttunen, J., Pekkonen, E., Kivisaari, R., Autti, T. & Kähkönen, S. (2008) Modulation of somatosensory evoked fields from SI and SII by acute GABA A-agonism and paired-pulse stimulation. *NeuroImage*, **40**, 427–434.
- Kujirai, T., Caramia, M.D., Rothwell, J.C., Day, B.L., Thompson, P.D., Ferbert, A., Wroe, S., Asselman, P. & Marsden, C.D. (1993) Corticocortical inhibition in human motor cortex. *J. Physiol.*, **471**, 501–519.
- Lenz, M., Höfken, O., Stude, P., Lissek, S., Schwenkreis, P., Reinersmann, A., Frettlöh, J., Richter, H., Tegenthoff, M. & Maier, C. (2011) Bilateral somatosensory cortex disinhibition in complex regional pain syndrome type I. *Neurology*, **77**, 1096–1101.
- Lenz, M., Tegenthoff, M., Kohlhaas, K., Stude, P., Höfken, O., Gatica Tossi, M.A., Kalisch, T., Kowalewski, R. & Dinse, H.R. (2012) Increased excitability of somatosensory cortex in aged humans is associated with impaired tactile acuity. *J. Neurosci.*, **32**, 1811–1816.
- McCarthy, G., Wood, C.C. & Allison, T. (1991) Cortical somatosensory evoked potentials. I. Recordings in the monkey *Macaca fascicularis*. *J. Neurophysiol.*, **66**, 53–63.
- McLaughlin, D.F. & Kelly, E.F. (1993) Evoked potentials as indices of adaptation in the somatosensory system in humans: a review and prospectus. *Brain Res. Brain Res. Rev.*, **18**, 151–206.
- Mitzdorf, U. (1985) Current source-density method and application in cat cerebral cortex: investigation of evoked potentials and EEG phenomena. *Physiol. Rev.*, **65**, 37–100.
- Musselwhite, M.J. & Jeffreys, D.A. (1983) Visual evoked potentials to double-pulse pattern presentation. *Vision Res.*, **23**, 135–143.
- Nicholson Peterson, N., Schroeder, C.E. & Arezzo, J.C. (1995) Neural generators of early cortical somatosensory evoked potentials in the awake monkey. *Electroen. Clin. Neuro.*, **96**, 248–260.
- Percaccio, C.R., Engineer, N.D., Pruetto, A.L., Pandya, P.K., Moucha, R., Rathbun, D.L. & Kilgard, M.P. (2005) Environmental enrichment increases paired-pulse depression in rat auditory cortex. *J. Neurophysiol.*, **94**, 3590–3600.
- Porter, J.T. & Nieves, D. (2004) Presynaptic GABAB receptors modulate thalamic excitation of inhibitory and excitatory neurons in the mouse barrel cortex. *J. Neurophysiol.*, **92**, 2762–2770.
- Ragert, P., Becker, M., Tegenthoff, M., Pleger, B. & Dinse, H.R. (2004) Sustained increase of somatosensory cortex excitability by 5 Hz repetitive transcranial magnetic stimulation studied by paired median nerve stimulation in humans. *Neurosci. Lett.*, **356**, 91–94.
- Schwartz, M. & Shagass, C. (1964) Recovery functions of human somatosensory and visual evoked potentials. *Ann. N. Y. Acad. Sci.*, **112**, 510–525.
- Shagass, C. & Schwartz, M. (1964) Recovery functions of somatosensory peripheral nerve and cerebral evoked responses in man. *Electroen. Clin. Neuro.*, **17**, 126–135.
- Takahashi, T., Forsythe, I.D., Tsujimoto, T., Barnes-Davies, M. & Onodera, K. (1996) Presynaptic calcium current modulation by a metabotropic glutamate receptor. *Science*, **274**, 594–597.
- Urbano, A., Babiloni, F., Babiloni, C., Ambrosini, A., Onorati, P. & Rossini, P.M. (1997) Human short latency cortical responses to somatosensory stimulation. A high resolution EEG study. *NeuroReport*, **8**, 3239–3243.
- Vaughan, J.H. & Arezzo, J. (1988) The neural basis of event-related potentials. In Picton, T. (Ed.), *Human Event-Related Potentials*. Elsevier, Amsterdam, pp. 45–96.
- Wehr, M. & Zador, A.M. (2005) Synaptic mechanisms of forward suppression in rat auditory cortex. *Neuron*, **47**, 437–445.
- Werhahn, K.J., Kunesch, E., Noachtar, S., Benecke, R. & Classen, J. (1999) Differential effects on motorcortical inhibition induced by blockade of GABA uptake in humans. *J. Physiol.*, **517**, 591–597.
- Wolters, A., Schmidt, A., Schramm, A., Zeller, D., Naumann, M., Kunesch, E., Benecke, R., Reiners, K. & Classen, J. (2005) Timing-dependent plasticity in human primary somatosensory cortex. *J. Physiol.*, **565**, 1039–1052.
- Ziemann, U., Lönnecker, S., Steinhoff, B.J. & Paulus, W. (1996) The effect of lorazepam on the motor cortical excitability in man. *Exp. Brain Res.*, **109**, 127–135.
- Ziemann, U., Muellbacher, W., Hallett, M. & Cohen, L.G. (2001) Modulation of practice-dependent plasticity in human motor cortex. *Brain*, **124**, 1171–1181.