

plate, Cdk5 and p35 (a neuronal-specific activator of Cdk5) are required. Both molecules are present in migrating neurons, and inactivation of either of them produces a phenotype somewhat similar to reeler in that cells are positioned in an outside-in gradient<sup>6,7</sup>. However, by contrast to the adult reeler, in Cdk5/p35 mutants there is a relatively normal layer I below the subplate layer<sup>8,9</sup>, suggesting that in this mutant cortical plate cells respond normally to Reelin.

An interesting observation is that in the Cdk5<sup>-/-</sup> mutant, the subplate remains within the cortical plate during development; that is, early-born cortical plate cells are able to traverse the subplate, but late-born cells become positioned below the subplate, forming an underplate. This does not occur in the p35<sup>-/-</sup> mutant, in which all or most cortical plate cells are able to cross the subplate<sup>8</sup>. It is possible that late-born cortical plate cells require Cdk5 to cross the subplate, but are activated by a factor other than p35 (perhaps p39)<sup>10</sup>. An additional point is that the defect in p35<sup>-/-</sup> mutants is more severe in dorsomedial cortex, but lateral and ventrolateral cortices are less affected by the mutation, again suggesting that in the latter regions there might be other factors, in addition to p35, that activate Cdk5.

Based on these observations, I proposed an hypothesis suggesting that Cdk5 was a key factor in the evolutionary origin of the isocortical inside-out gradient<sup>5</sup>. This hypothesis postulated that in the cortical plate of the Cdk5/p35 mutant, and perhaps in that of an ancestral mammal-like reptile with an outside-in pattern of migration, there is, or was, a putative factor that blocks the migration of cells past previously formed cortical-plate cells. Because of the role of Cdk5 in promoting neurite outgrowth and modifying the cytoskeleton<sup>11-13</sup>, and the fact that Cdk5/p35 are expressed in migrating neurons, I suggested that the Cdk5/p35 pathway allows migrating cells to bypass this migratory-suppressing factor. Recently, Cdk5/p35 have been found to be associated with a  $\beta$ -catenin-N-cadherin in the cerebral cortex<sup>14</sup>. This study showed that inhibition of Cdk5 kinase activity facilitates *in vitro* N-cadherin mediated neuronal aggregation. Similarly, p35<sup>-/-</sup> cells also aggregate to a greater degree compared with wild-type neurons. These findings suggest that, *in vivo*, Cdk5 avoids the arrest of neuronal

migration below older cells by inhibiting N-cadherin-dependent cell aggregation.

The Reelin pathway might have participated in the origin of the isocortex by acting as a barrier for cellular migration into the marginal zone<sup>5</sup>, and perhaps by contributing to the organized cellular array in the cortical plate, which facilitates the migration of cells past previously formed layers. This hypothesis is consistent with the fact, mentioned by Bar *et al.*<sup>1</sup>, that the *reelin* gene is faintly expressed in the reptilian and avian cortical marginal zones, in spite of an otherwise evolutionarily conserved canvas of expression in other brain regions. In these animals, relatively low levels of Reelin might be sufficient to arrest the migration of the few cells that reach the marginal zone. Alternatively, a signal that is different from Reelin could prevent migration into the marginal zone of birds and reptiles. In the mammalian hippocampus and dentate gyrus, the reeler mutation leaves the marginal zone relatively intact<sup>4</sup>, suggesting that other factors, in addition to Reelin, are acting as stop signals in these zones. Perhaps a similar situation occurs in reptiles and birds.

#### Francisco Aboitiz

Programa de Morfología, ICBM, Facultad de Medicina, U. de Chile, P.O. Box 70079, Santiago 7, Chile.

e-mail: faboitiz@machi.med.uchile.cl

#### References

- 1 Bar, I. *et al.* (2000) The evolution of cortical development. An hypothesis based on the role of the Reelin signaling pathway. *Trends Neurosci.* 23, 633-638
- 2 Frotscher, M. (1998) Cajal-Retzius cells, reelin and the formation of layers. *Curr. Opin. Neurobiol.* 8, 570-575
- 3 Dulabon, L. *et al.* (2000) Reelin binds  $\alpha 3 \beta 1$  integrin and inhibits neuronal migration. *Neuron* 27, 33-44
- 4 Caviness, V.S. *et al.* (1988) The reeler malformation. Implications for neocortical histogenesis. In *Cerebral Cortex Vol. 7. Development and Maturation of the Cerebral Cortex* (Peters, A. and Jones, E.G., eds), New York, Plenum Press
- 5 Aboitiz, F. (1999) Evolution of isocortical organization. A tentative scenario including roles of reelin, p35/cdk5 and the subplate zone. *Cereb. Cortex* 9, 655-661
- 6 Oshima, T. *et al.* (1996) Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc. Natl. Acad. Sci. U. S. A.* 93, 11173-11178
- 7 Chae, T. *et al.* (1997) Mice lacking p35, a neuronal specific activator of cdk5, display cortical

lamination defects, seizures, and adult lethality. *Neuron* 18, 29-42

- 8 Gilmore, E.C. *et al.* (1998) Cyclin-dependent kinase 5-deficient mice demonstrate novel developmental arrest in cerebral cortex. *J. Neurosci.* 18, 6370-6377
- 9 Kwon, Y.T. and Tsai, L.H. (1998) A novel disruption of cortical development in p35(-/-) mice distinct from reeler. *J. Comp. Neurol.* 395, 510-522
- 10 Walsh, C.A. and Goffinet, A.M. (2000) Potential mechanism of mutations that affect neuronal migration in man and mouse. *Curr. Opin. Genet. Dev.* 10, 270-274
- 11 Paglini, G. *et al.* (1998) Evidence for the participation of the neuron-specific cdk5 activator p35 during laminin-enhanced axonal growth. *J. Neurosci.* 18, 9858-9869
- 12 Kwon, Y.T. *et al.* (1999) Callosal axon guidance defects in p35(-/-) mice. *J. Comp. Neurol.* 415, 218-229
- 13 Nikolic, M. *et al.* (1998) The p35/cdk5 kinase is a neuron-specific Rac effector that inhibits Pak1 activity. *Nature* 395, 194-198
- 14 Kwon, Y.T. *et al.* (2000) Regulation of N-cadherin-mediated adhesion by the p35-Cdk5 kinase. *Curr. Biol.* 10, 363-372

## Time-variant processing in V1: from microscopic (single cell) to mesoscopic (population) levels

In their article, Wörgötter and Eysel<sup>1</sup> summarize recent trends on the understanding of receptive field (RF) dynamics and on the contextual and attentional influences on RF properties. They correctly point out that cortical RFs are highly dynamic entities embracing more than the sum of the full spatial and temporal response properties of single cells.

Given that this endeavour requires intricate interactions between large numbers of cells, the question arises of how the outcome of this cooperativity can be conceptualized and analyzed at the mesoscopic level of neural populations. From a phenomenological point of view the idea to analyze entire populations is a rather inescapable consequence of the observation that huge numbers of neurons are activated, even after the simplest form of stimulation<sup>2-4</sup>. We recently introduced a new approach to study population activity in early sensory cortices. Our goal was to investigate cortical activity distributions

in the coordinates of the respective stimulus space in order to explore cooperative processes<sup>5–8</sup>.

To explore the contribution of many cells to a common stimulus representation across visual space, we constructed dynamic ‘population-RFs’<sup>6</sup>. A Gaussian interpolation procedure (for technical details see Refs 8,9) was used to depict the entire temporal structure of neural responses of nearly 200 single cells recorded in area 17. The population dynamics were captured using the time-slice technique, which is frequently used for calculation of single-cell dynamic RFs (time-slicing and reverse correlation are identical assuming the use of simple on–off stimuli). However, the spatio-temporal evolution of ‘population-RFs’ differed compared with single cell RFs. First, the time-scale of population RFs was much shorter: population activity was characterized by a gradual build-up and decay lasting only 40–50 ms, whereas single-cell dynamics typically vary over much longer periods<sup>10–16</sup>. Second, the profile of the population RFs indicated a remarkable spatio-temporal coherence across visual space (Fig. 1).

To understand these differences one has to recall how the construction of population RFs was achieved: because our main goal was to study the population response to a common stimulus (i.e. the contributions of all neurons responding to the given stimulus), its position was kept fixed irrespective of the location or specific properties of each RF analyzed. This procedure becomes intuitively clear when taking into account the fact that under natural viewing conditions, stimuli are similarly distributed in arbitrary ways across RFs. What is the impact of this procedure?

Using the common-response plane technique for assessment of RF dynamics, different positions within the RF of a cell are stimulated: when the time course of each position reveals substantial and systematic differences, the cell is said to be spatio-temporally non-separable. Assuming that about 50% of V1 neurons are non-separable and that our population response is a representative sample of V1, we would expect that separable and non-separable cells contribute in approximately equal proportions. However, the complex spatio-temporal behavior that is characteristic for non-




Fig. 1. Time courses of population activity in response to squares of light (0.4°) flashed at different locations in the visual field. Two-dimensional distributions of population activity were obtained from extracellular recordings of 178 single neurons in cat primary visual cortex. The activity distributions were calculated for consecutive time windows of 10 ms duration covering the period from 30 to 80 ms after stimulus onset. The frame with the cross-hair illustrates the analyzed portion of the visual field (2.8 × 2.0°). The positions of the stimuli are indicated by the small square outlined in white within the stimulus frame. The spatial scale of the stimuli and the stimulus separations matches the configurations used for assessment of non-separability in single cell approaches. Note the similarities in time course and the coherence of activation across the entire population. Reproduced, with permission, from Ref. 6.

separable cells appeared not to be preserved in our population dynamics. We therefore argue that, at the level of populations, each position in the visual field is well-represented by a large pool of neurons that respond maximally to any potential stimulus configuration. Consequently, the structure of single-cell RF dynamics, characterized by highly complicated distributions of activity in space and time, transforms into much simpler spatio-temporal activity patterns (Fig. 1). A comparable coherence of overall response dynamics has also been observed for somatosensory cortical populations<sup>17</sup>.

The transformation of the diversified spatio-temporal behavior of single cells into coherent ensemble dynamics might indicate a qualitative step of time-variant processing from microscopic into mesoscopic (e.g. population) levels<sup>18,19</sup>, in which the aspect of non-separability might become obsolete.

Such a lack of separability and complexity in response to stimuli that are very simple in shape and timing, however, does not necessarily imply that the single-cell dynamics are meaningless. Indeed, we assume that the detailed, idiosyncratic time structure is a consequence of cooperative processes that, in turn, are generated by non-linear time-dependent interaction effects across the entire neural network. At the population level, consistent coherent dynamics become observable that allow a direct link

between physiological and psychophysical phenomena. Conceivably, the utilization of more-complicated and demanding spatio-temporal stimulus configurations might allow a better understanding of cortical function. In fact, recent evidence from our laboratory indicates that moving stimuli are represented as coherently moving waves of population activity and that these waves emerge from active processes underlying lateral pre-activation and integration across the entire population<sup>7,20</sup>. Interestingly, the representations of moving stimuli were characterized by significantly shorter latencies compared with representations of a flash (D. Jancke *et al.* unpublished observations). Psychophysically, a flashed and a moving stimulus presented at the same location are perceived as being displaced<sup>21</sup>. It is therefore tempting to speculate that the latency differences observed for populations contribute to the so-called ‘flash-lag’ effect.

**Hubert R. Dinse\***

Inst. Neuroinformatics, Dept Theoret.  
Biol. Ruhr-Univ. Bochum. ND 04,  
D-44780 Bochum, Germany.  
\*e-mail: hubert.dinse@  
neuroinformatik.ruhr-uni-bochum.de

**Dirk Jancke**

Weizmann Institute of Science, Dept of  
Neurobiology, P.O. Box 26, 76100, Rehovot,  
Israel.

## References

- 1 Wörgötter, F. and Eysel, U. (2000) Context, state and the receptive fields of striate cortex cells. *Trends Neurosci.* 23, 497–503
- 2 Grinvald, A. *et al.* (1994) Cortical point-spread function and long-range lateral interactions revealed by real-time optical imaging of macaque monkey primary visual cortex. *J. Neurosci.* 14, 2545–2568
- 3 Godde, B. *et al.* (1995) Optical imaging of rat somatosensory cortex reveals representational overlap as topographic principle. *NeuroReport* 7, 24–28
- 4 Chen-Bee, C.H. and Frostig, R.D. (1996) Variability and interhemispheric asymmetry of single-whisker functional representations in rat barrel cortex. *J. Neurophysiol.* 76, 884–894
- 5 Dinse, H.R. *et al.* (1996) Dynamics of population representations of visual and somatosensory cortex based on spatio-temporal stimulation. In *Progress in Neural Information Processing. International Conference on Neural Information Processing. ICONIP'96* (Amari, S. *et al.*, eds), pp. 1285–1290, Springer
- 6 Jancke, D. *et al.* (1999) Parametric population representation of retinal location: neuronal interaction dynamics in cat primary visual cortex. *J. Neurosci.* 19, 9016–9028
- 7 Jancke, D. (2000) Orientation formed by a spot's trajectory: a two-dimensional population approach in primary visual cortex. *J. Neurosci.* 20, RC86
- 8 Dinse, H.R. and Jancke, D. (2001) Comparative population analysis of cortical representations in parametric spaces of visual field and skin: a unifying role for nonlinear interactions as a basis for active information processing across modalities. In *Advances in Population Coding* (Nicolleis, M.A.L., ed.) Progress in Brain Research. Elsevier, in press
- 9 Erlhagen, W. *et al.* (1999) The distribution of neuronal population activation (DPA) as a tool to study interaction and integration in cortical representations. *J. Neurosci. Methods* 94, 53–66
- 10 Jones, J.P. and Palmer, L.A. (1987) The two-dimensional spatial structure of simple receptive fields in cat striate cortex. *J. Neurophysiol.* 58, 1187–1211
- 11 Dinse, H.R. *et al.* (1990) A temporal structure of cortical information processing. *Concepts Neurosci. (CINS)* 1, 199–238
- 12 Dinse, H.R. *et al.* (1991) Temporal aspects of cortical information processing – cortical architecture, oscillations, and non-separability of receptive field organization. In *Neuronal Cooperativity – Models and Experiments* (Krüger, J., ed.), pp. 68–104, Springer
- 13 Shevelev, I.A. *et al.* (1992) Dynamics of responses of V1 neurons evoked by stimulation of different zones of receptive field. *Neuroscience* 51, 445–450
- 14 DeAngelis, G.C. *et al.* (1993) Spatiotemporal organization of simple-cell receptive fields in the cat's striate cortex. I. General characteristics and postnatal development. *J. Neurophysiol.* 69, 1091–1117
- 15 Eckhorn, R. *et al.* (1993) The RF-cinematogram. A cross-correlation technique for mapping several visual receptive fields at once. *Biol. Cybern.* 69, 37–55
- 16 Dinse, H.R. (1994) A time-based approach towards cortical functions: Neural mechanisms underlying dynamic aspects of information processing before and after postontogenetic plastic processes. *Physica D* 75, 129–150
- 17 Kalt, T. *et al.* (1996) Dynamic population coding in rat somatosensory cortex. *Soc. Neurosci. Abstr.* 22, 105
- 18 Freeman, W.J. (1999) *How Brains Make Up Their Minds*. Weidenfeld & Nicolson General
- 19 Freeman, W. *et al.* (2000) Analysis of spatial patterns of phase in neocortical gamma EEGs in rabbit. *J. Neurophysiol.* 84, 1266–1278
- 20 Jancke, D. *et al.* (1996) Reconstruction of motion trajectories from the dynamic population representation of neurons in cat visual cortex. *Soc. Neurosci. Abstr.* 22, 646
- 21 MacKay, D.M. (1958) Perceptual stability of a stroboscopically lit visual field containing self-luminous objects. *Nature* 181, 507–508

## Glia-neuronal culture models – do we need to change the paradigms?

The review by Fields and Stevens<sup>1</sup> was welcome in its much needed emphasis on the roles of the neuroglia in the CNS and PNS, in this case focusing on how the important extracellular signaling molecule, ATP, might direct glia–neuron interactions. Several studies suggesting intriguing possibilities were described. However, a major deficiency that is apparent from this review is that almost all the studies, so far, have been carried out in various culture or co-culture systems or easily accessible PNS systems. There is a reasonably abundant body of work indicating differences between astroglial and other CNS glial properties in culture and *in situ*, and also similarities. The problem with this situation is that the similarities or differences are unpredictable so one always needs *in situ* knowledge of the property in question. Similarities and differences are also probably a function of how probing the questions are. So what systems should be used to get the reliable information that is required to know that our ideas of how the PNS or CNS works will not be forever flawed because it is not known what the 'glia' are actually doing? Currently the most reliable systems are tissue slices and freshly isolated cells (which provide information on the individual properties without the changes in gene expression that occur as a result of the culture environment), and of course the classical histological techniques with all their remarkable up-dates. Another feature

that is well known but needs to be considered more rigorously is that glia refers to three major classes of different cells in the CNS with further subtypes, which in turn might shelter an unknown number of functional variants.

Resolution of these issues only appears possible by a shift of emphasis of a critical mass of productive laboratories from neuron to glial studies. This could be instituted by targeting funds to the glial field, but at whose behest and with what justification would we violate the principle of investigator-initiated science with its demonstrated creativity? To paraphrase Newton\* we still support the view that it is not for one group to direct the studies of another investigator. However, we all know that peer review, the litmus test of scientific merit, in fact does this all the time, but in a reactive rather than proactive manner. The smallness of the 'glial' field and that technically easy, rather than harder and more reliable, experiments are mainly carried out, are both the cause and effect of the size of the field and the low amounts of reliable data. So when the city neuron neuroscientists ask their poor country glial colleagues 'so what do glia actually do?', all we can say is 'we're working on it as best we know how. We'll have some answers for you as soon as we can.'

Harold K. Kimelberg

Albany Medical College, Albany, NY 12208, USA.

e-mail: kimelbH@mail.amc.edu

### Reference

- 1 Fields, R.D. and Stevens, B. (2000) ATP: an extracellular signaling molecule between neurons and glia. *Trends Neurosci.* 23, 625–633

\*"But he knows well yt it is not for one man to prescribe Rules to ye studies of another." From 1672 letter by Newton to Henry Oldenburg, secretary to the Royal Society, concerning the despised Robert Hooke.

## Response: glia-neuronal culture models – do we need to change the paradigms?

Kimelberg<sup>1</sup> raises several important points concerning studies of purinergic receptors in glia. Sensitivity to the importance of potential differences